



RTB

National plan for improving seed potato in Georgia

M A R C H 2 0 1 9

Authors

R. Mdivani, J. Andrade, I. Girard,
R. Shavgulidze, N. Khatiashvili,
G. Forbes (senior author).



RESEARCH
PROGRAM ON
Roots, Tubers
and Bananas



CIP
INTERNATIONAL
POTATO CENTER

Funded by the Republic of Austria





RTB

National plan for improving seed potato in Georgia

M A R C H 2 0 1 9

Authors

R. Mdivani, J. Andrade, I. Girard,
R. Shavgulidze, N. Khatishvili,
G. Forbes (senior author).



RESEARCH
PROGRAM ON
Roots, Tubers
and Bananas



CIP
INTERNATIONAL
POTATO CENTER

Funded by the Republic of Austria



Correct citation:

CIP (2019). Plan for improving seed potato in Georgia. RTB Working Paper. CIP, Lima, Peru

Published by the CGIAR Research Program on Roots, Tubers and Bananas

The CGIAR Research Program on Roots, Tubers and Bananas (RTB) is a partnership collaboration led by the International Potato Center implemented jointly with Bioversity International, the International Center for Tropical Agriculture (CIAT), the International Institute of Tropical Agriculture (IITA), and the Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), that includes a growing number of research and development partners. RTB brings together research on its mandate crops: bananas and plantains, cassava, potato, sweetpotato, yams, and minor roots and tubers, to improve nutrition and food security and foster greater gender equity especially among some of the world's poorest and most vulnerable populations.

Contact:

RTB Program Management Unit
International Potato Center (CIP)
Apartado 1558, Lima 12, Peru
rtb@cgiar.org • www.rtb.cgiar.org

CIP-Georgia office
6, Marshal Gelovani Ave. 0159, Tbilisi, Georgia
r.mdivani@cgiar.org; www.cipotato.org

CIP thanks all donors and organizations which globally support its work through their contributions to the CGIAR Trust Fund.
<https://www.cgiar.org/funders/>

© International Potato Center on behalf of RTB



This working paper is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/4.0/>.

Disclaimer:

This RTB Working Paper is intended to disseminate research and practices about production and utilization of roots, tubers and bananas and to encourage debate and exchange of ideas. The views expressed in the papers are those of the author(s) and do not necessarily reflect the official position of RTB, CGIAR or the publishing institution.

Contents

Acronyms	5
Introduction.....	6
Objectives and approach.....	7
Methodology	7
Beneficiaries	7
Potencial benefits.....	8
Governance	8
Funding.....	8
Resources	9
Components and activities	9
Component 1: Adoption of varieties that degenerate slowly due to their natural resistance to the organisms that cause PSD	9
Activity 1.1. Import new genotypes from CIP and other sources.....	9
Activity 1.2. Identify and catalogue virus resistance in candidate varieties.....	9
Activity 1.3. Participatory varietal trials	10
Activity 1.4. Release and dissemination of new varieties	10
Activity 1.5. Adoption studies of new varieties	10
Component 2: On-farm practices to maintain or even improve seed quality.....	10
Activity 2.1. Adaptation of techniques and training materials.....	10
Activity 2.2. Training of extension agents and farmers.....	10
Activity 2.3. Monitoring seed health	11
Component 3: Strategic use of certified seed	11
Activity 3.1. Market chain survey	11
Activity 3.2. Analysis to develop joint business opportunities	11
Activity 3.3. Implementation of joint innovations.....	12
References.....	12

Acronyms

ADA: Austrian Development Agency

APSIM: Agricultural Production System Simulator

CIP: International Potato Center (*Centro Internacional de la Papa*)

ELISA: Enzyme-linked immunosorbent assay

FAO: Food and Agriculture Organization of the United Nations

FESC: Farmers' Extension Service Centers

GEL: Georgian Lari

GLC: Grain Logistic Company

ISH: Integrated seed health

MEPA: Ministry of Environmental Protection and Agriculture

MoA: Ministry of Agriculture

PLRV: Potato leafroll virus

PMCA: Participatory market chain approach

PS: Positive selection

PSD: Potato seed degeneration

PVY: Potato virus Y

R&D: Research and development

RTB: CGIAR Research Program on Roots, Tubers and Bananas

SPP: Seed potato platform

SRCA: Scientific Research Center of Agriculture

SSA: Sub-Saharan Africa

Abstract

Keywords: potato, seed potato, Georgia, national plan

Acknowledgements

This research was undertaken as part of, and funded by, the CGIAR Research Program on Roots, Tubers and Bananas and the International Potato Center supported by [CGIAR Trust Fund contributors](#). Funding support for this work was provided by the Austrian Development Agency (ADA) through the project *Enhancing Rural Livelihoods in Georgia: Introducing Integrated Seed Health Approaches to Local Potato Seed Systems*.

Introduction

Potato is an important crop in Georgia, with approximately 20,000 ha under production, and annual per capita consumption of 55 kg (Geostat 2019). Nearly all potato is consumed fresh, and Georgians consider the crop as their ‘second bread’.

Despite potato’s importance in the diet and culture, yields remain low: at roughly 11 t/ha, the national average is far below yields of 40 t/ha or more found in many northern European countries. Research has demonstrated that with current varieties, attainable yields (on experiment stations and by the best, high-input farmers) are 25–30 t/ha (Carli et al. 2010; Mdivani 2012). The proposed Seed Plan, described in this report, will help reduce the large gap between actual and attainable yields.

Tubers serve as the planting material, or seed, of the potato. The poor quality of seed potato is one of the main causes of the low productivity and reduced profitability of Georgia’s potato crop (Carli et al. 2010; MoA 2015). As a vegetatively propagated crop, seed potato accumulates pests and pathogens—particularly viruses—over successive seasons of cultivation. This leads to a reduction in yield capacity and/or quality of the product, a phenomenon referred to as *potato seed degeneration* (PSD) (Thomas-Sharma et al. 2015).

In response to the problem of PSD, seed certification systems in the United States and Europe were developed in the early part of the 20th century (Frost et al. 2013). These systems have been highly successful, making high-quality seed readily available to farmers and virtually eliminating PSD in high-income countries. Many efforts to establish certified seed systems for potato in low-income economies, however, have had little success; most potato farmers in these areas

source seed from the informal seed system (e.g. produced on-farm, acquired from neighbors or local markets) (McGuire and Sperling 2016). Such seed can often be highly degenerated and cause major reductions in productivity (Thomas-Sharma et al. 2015).

Currently, most potato seed in Georgia is self-saved (as part of ware production, i.e. potatoes for human consumption), or obtained from local sources, by small-scale farmers who are unaware of practices to maintain seed quality (e.g. proper storage). Most of this seed is severely affected by PSD (MoA 2015). Even the portion of seed that is imported and has been multiplied in country is sold without certification or proper bagging and labeling. Farmers routinely do not use on-farm practices aimed at maintaining seed quality, seed treatment, and seed selection. Nor have they had systematic, formal access to training that would enable them to properly manage their potato harvests (Carli et al. 2010).

Direct access to imported seed is limited in Georgia, and such seed is very expensive due to the high relative value of the Euro currency. Seed multiplied from imported seed costs 2–3 GEL¹/kg, whereas seed from local sources costs 0.5–0.8 GEL/kg. That the quality of imported seed is not guaranteed is another issue; for example, seed quality is sometimes greatly diminished by poor transport conditions. In addition, seed imported from Europe is of varieties that generally lack resistance to pathogens causing PSD, principally the two most yield-reducing viruses, potato virus Y (PVY) and potato leafroll virus (PLRV). We are unaware of studies demonstrating the susceptibility of potato varieties currently imported into Georgia, but many of the same ones were found to be highly susceptible in recent studies done in Tajikistan (CIP 2016). All these factors have caused the majority of the seed potato planted in Georgia to be highly affected by PSD (MoA 2015).

Georgia has excellent agronomic and environmental conditions in highland areas where high-quality seed can easily be produced (Carli et al. 2010). During the Soviet era, however, both seed production and seed certification management were centralized in the Soviet Union and basic seed was produced in Russia, Belarus, and Ukraine. This seed was then shipped to other Soviet countries, including Georgia, for further multiplication and eventual certification. Thus, a full seed production chain did not exist in Georgia at that time. With the collapse of the Soviet Union, the flow of basic seed and support for certification stopped, effectively ending certified seed production in Georgia. Subsequently, some high-quality seed has been imported from Europe, but in small amounts; even after multiplication in Georgia it represented less than 10% of seed needs (Geostat). There is currently no control on the quality of imported seed.

¹ Georgian Lari (GEL). Conversion rate in March 2019 is approximately USD 1 = GEL 2.7

Objectives and approach

The objective of the Plan for Improving Seed Potato in Georgia (hereafter referred to as Seed Plan) is to provide direction to appropriate Georgian authorities for addressing constraints related to quality seed production in Georgia, especially PSD, in the next five years, 2019-2024. This plan is based on the *Integrated Seed Health* (ISH) approach which describes three tactics to manage PSD (Thomas-Sharma et al. 2015): the adoption of varieties that degenerate slowly due to their natural resistance to the organisms that cause PSD; on-farm practices to maintain or even improve seed quality at the farm level; and strategic use of certified seed (i.e. buying in quantities and at frequencies that correspond to farmers' needs). Each of these tactics corresponds to a component in the current Seed Plan.

Important. This Seed Plan should be closely linked to a plan to develop the potato value chain in Georgia. This is because potato yields are expected to increase as a result of the activities implemented in the Seed Plan, and this surplus must be taken up by increased local consumption, or international markets. If not, potato prices could collapse and potato farmers' economies would be damaged.

Methodology

A first draft of the Seed Plan was developed and written by International Potato Center (CIP) experts in the frame of the Austrian Development Agency (ADA) funded project *Enhancing Rural Livelihoods in Georgia: Introducing Integrated Seed Health Approaches to Local Potato Seed Systems*. One of the expected outputs of the ADA project is to have a potato seed plan incorporating integrated seed health (ISH) strategies developed for Georgia.

The Seed Plan was improved using information from a baseline study on potato seed systems conducted by CIP in the second half of 2017. Based on collected information, CIP mapped existing seed systems using network analysis, existing information, and expert consultation, and from this developed seed degeneration scenarios with modeling. CIP elaborated the role of seed quality in yield gaps using the Agricultural Production System Simulator (APSIM) model (Holzworth et al. 2014). Based on the outputs of the model, CIP developed a draft Seed Plan. In September 2018, CIP organized a stakeholder workshop at which the plan was presented. Stakeholders provided feedback to the draft Seed Plan and, as a result, CIP incorporated new elements in a revised version, which will be formally presented to the Ministry of Agriculture in late March 2019.

Beneficiaries

The target beneficiaries of the activities implemented as part of the Seed Plan are potato farmers in Georgia, who will benefit from improved quality of planting material, which will boost their productivity and yield stability, and consequently increase their families' income and quality of life. Increased profitability of the potato crop could also benefit consumers through lower and/or more stable prices. An improved seed system will also lead to more efficient diffusion of new varieties, which has benefits to farmers and consumers that transcend productivity.

Potential benefits

Although it is difficult to predict quantitative benefits to the target group, previous experiences of CIP researchers would indicate that very significant increases in productivity can be expected by improving seed quality. For example, researchers in Sub-Saharan Africa (SSA) have shown yield increases of around 30% on average with one season of positive selection (PS) (Schulte-Geldermann et al. 2012). PS is not 100% efficient—farmers might miss latently or even mildly infected plants—so some yield loss would still occur. Presumably, these studies present a highly conservative estimate of actual yield losses in farmers' fields caused by degeneration. Also, raising awareness of how to manage properly potato fields would facilitate the adoption of good agronomic practices, further increasing current potato yields in Georgia. There is also strong evidence that host resistance to PSD pathogens benefits farmers. Several seed degeneration studies have compared two or more varieties, sometimes with known levels of resistance, and indicated differences in yield loss. In Kenya for example, after four seasons the overall yield reduction in genotypes resistant to multiple viruses ranged from 5% to 33%, whereas yield reduction in the local Ugandan and Kenyan varieties ranged from 56% to 58% (Ondity et al. 2013).

Governance

The implementation of the Seed Plan will be led by the Ministry of Agriculture (MoA). A multi-stakeholder platform (see example in Thiele et al. 2011), or Seed Potato Platform (SPP), will be created to support the implementation of the plan by creating a common agenda for stakeholders. The SPP will include stakeholders related directly to seed production, distribution, and use, such as national research and development (R&D) organizations², farmers' cooperatives, seed importers, seed producers, donors, and international R&D organizations (CIP, FAO).

² Ministry of Environmental Protection and Agriculture (MEPA); Scientific Research Center of Agriculture (SRCA); MoA—Farmers' Extension Service Centers (FESC); MoA—Grain Logistic Company (GLC)

Stakeholders will be identified and mapped through a survey conducted in individual interviews and focus group discussions. Other potential stakeholders will be identified using criteria such as location of farms and farmed area. Roles and responsibilities of all stakeholders in the SPP will be defined during the survey exercise.

The SPP should be part of a wider platform to develop the potato value chain in Georgia, which will include the above stakeholders, plus others related to markets and consumption, such as processors, supermarkets, and consumer associations.

Funding

The design of this plan is supported by the project *Enhancing Rural Livelihoods in Georgia: Introducing Integrated Seed Health Approaches to Local Potato Seed Systems* funded by the Austrian Development Agency (ADA) (CIP 2018). Likewise, several activities described below are currently being funded and implemented by this project. The MoA and other partners are expected to fund significant portions of this plan.

Resources

CIP is working closely with the MoA and other partners to provide support on the design and implementation of this plan. A comprehensive set of bibliographical resources, including research papers, training materials, and websites is accessible through the internet (see References).

Components and activities

Below, the details of the plan are provided for the three components of Integrated Seed Health. Each component has several activities and for each we describe the methodology, the expected outputs and timeframe, ongoing activities implemented by the ADA-funded project, and the action plan (to be developed during the start-up workshop) (Excel document, attached).

Component 1: Adoption of varieties that degenerate slowly due to their natural resistance to the organisms that cause PSD

The activities in this component are aimed at improving the adoption of varieties resistant to viruses, the main cause of PSD in Georgia and elsewhere. However, these varieties must first be developed, because there is indirect evidence (Forbes et al. 2018) that shows that most of the varieties grown in Georgia are susceptible to viruses. In addition, virus resistance is just one of the traits considered when developing a new variety. Therefore, this component includes activities that are part of a variety development program, but that can be included in the current Seed Plan (to be discussed during the start-up workshop of the implementation process).

Activity 1.1. Import new genotypes from CIP and other sources. Potato experts (scientists, extension agents, processors, and farmers), in coordination with breeders from CIP and other organizations, will select potato genotypes from existing catalogues (e.g. CIP n.d.), as part of a variety development program. The expected output is 20 to 25 new potato genotypes available in Georgia every two or three years. This activity is well underway as part of the current ADA-funded project. A large number of new varieties is being evaluated in experimental trials on-station and a small number of candidate varieties with superior traits will be assessed in farmers' fields (see Activity 1.3). The Seed Plan could complement these existing efforts by identifying a mechanism for continued introduction and evaluation of new experimental potato genotypes.

Activity 1.2. Identify and catalogue virus resistance in candidate varieties. Between 20 and 25 potato genotypes will be planted in areas and seasons with high aphid populations and virus incidence to evaluate resistance to the main viruses (PVY and PLRV) following the protocol developed by CIP (CIP 2006). Evaluation for other biotic and abiotic constraints could also be included. Selection of areas will be based on agro-ecological conditions, as well as on availability of machinery, irrigation, and other factors. The expected output is a catalogue of potato genotypes characterized by their resistance to virus. In addition, the methodology for testing virus resistance is expected to be up taken by Georgian breeders to be implemented in a variety development program. Currently, the ADA-funded project is finishing the first cycle of evaluations of 32 potato genotypes.

Activity 1.3. Participatory varietal trials. Between 20 and 25 potato genotypes will be selected under field conditions by farmers, consumers, and scientists following a participatory methodology (De Haan et al. 2017) for three years, paying close attention to gender considerations (Mudege et al. 2017). Before planting, seeds will be sorted visually to prevent spread of pests and diseases. Also, pesticides will be used for seed treatment. Other cultural practices will be conducted

following local practices. The expected output is two to five potato genotypes, identified for local conditions, every two or three years, as part of a variety development program.

Activity 1.4. Release and dissemination of new varieties. Potato genotypes will be launched formally by Georgian authorities, following protocols developed in Component 3, as part of a variety development program. Dissemination of new varieties will include the production of certified seed (also addressed in Component 3) and distribution to farmers using several mechanisms, e.g. vouchers (Ogero et al. 2016) and small seed packs (FIPS n.d.). The expected output is two to five new potato varieties launched officially and in the hands of at least 5,000 farmers at the end of year five of this Plan.

Activity 1.5. Adoption studies of new varieties. Expert workshops will be used to monitor the degree of adoption of new varieties (see an example in Gatto et al. 2017), as part of a variety development program. The expected output is a study to identify the varieties with the highest adoption every five to seven years.

Component 2: On-farm practices to maintain or even improve seed quality

The activities in this component are aimed at improving farmers' capacities to manage their seed potato in the field, but these can also apply to seed multipliers. This component will be particularly important in the Southern Highlands and the Northern Highlands where an estimated 70% and 20% of seed, respectively, come from farmer-saved tubers (Forbes et al. 2018). It will also be important in the Midlands region to improve current use of seed imported from Europe by using seed replacement practices that combine the use of high-quality seed (Component 3) with on-farm practices. This component includes activities for adapting technologies and training materials to Georgian conditions, training extension agents and farmers, and monitoring progress on seed health.

Activity 2.1. Adaptation of techniques and training materials. Previously developed techniques and training materials on positive selection (Gildemacher et al. 2007a and 2007b), seed plot technique (Bryan 1983; Kinyua et al. 2015), replacement practices with certified seed (Obura et al. 2016), and techniques for managing seed-borne pests (e.g. late blight) (Caceres et al. 2008) will be adapted to Georgian conditions by experts and validated with farmers. Training resources will include not only printed materials, but also videos (see cassava example, Access Agriculture n.d.). The expected output is a set of techniques and training materials for improving farmer-saved seed that is adapted for Georgian conditions and available in the next two years, and then updated every five years.

Activity 2.2. Training of extension agents and farmers. Training of trainers' courses using previously developed materials (Gildemacher et al. 2007c) will be conducted to improve the capacities of extension agents to reach farmers and help them to manage their seed, using the techniques mentioned above. Methodologies to improve farmers' capacities will include farmer field schools (e.g. Wahyuning et al. 2006) and others, depending on the context. The intended output is to have 54 extension agents and 300 farmers trained on management practices for improving farmer-saved seed every year (the first cadre graduated after two years). Currently, the ADA-funded project has started training farmers on the positive selection technique.

Activity 2.3. Monitoring seed health. Although the baseline study provided valuable information on major seed-borne diseases (Forbes et al. 2018), it is important to identify new seed-borne pests and diseases, and monitor the progress of seed health as result of the introduction of new varieties and capacity development activities, described above. Seed tubers from a randomly selected sample of farmers will be collected and analyzed using ELISA tests or other diagnostic techniques. Depending on funding, we can include soil samples to analyze the incidence of soil-borne pathogens and nematodes. The expected output is a map with incidence and severity of major seed-borne potato pests and diseases, updated every two years. Currently, the ADA-funded project, using expert elicitation, has identified several seed-borne diseases, including potato wart, a quarantine disease caused by *Synchytrium endobioticum* (Forbes et al. 2018).

Component 3: Strategic use of certified seed

This component will promote the production of certified seed in Georgia (i.e. to develop the formal seed sector) and improve its use by implementing seed replacement practices that combine the use of high-quality seed and on-farm practices (Component 2). The activities will follow the *participatory market chain approach* (PMCA) (Bernet et al. 2006) and should be closely linked to activities aimed at developing the potato value chain in Georgia.

In the PMCA, stakeholders of the Seed Potato Platform (SPP) will interact to identify market opportunities and then develop innovations to take advantage of them. Innovations are categorized into three classes: commercial innovations (e.g. a new commercial product); technological innovations (e.g. a new agronomic technique); and institutional innovations (e.g. a new regulation or a new farmers' cooperative). The PMCA includes three phases: market chain survey; analysis to develop joint business opportunities; and implementation of joint innovations. Each of these phases corresponds to an activity in Component 3, which will be defined in detail by the stakeholders during the PMCA.

Activity 3.1. Market chain survey. The R&D organization leading the PMCA will conduct 20 to 40 interviews to identify key stakeholders and explore market opportunities and bottlenecks. The results will be presented at a public event where stakeholders will start interacting. The intended output is a report indicating key stakeholders, market opportunities, and bottlenecks. Examples of market opportunities include: a processing company that operates under a contract farming approach and needs to provide high-quality seed to its associated farmers; and demand for certified seed from a neighboring country. Examples of bottlenecks include lack of a technique for producing high-quality seed or for identifying key seed-borne pathogens; lack of a network of seed producers; and lack of a regulation to export seed. This activity will take approximately three months.

Activity 3.2. Analysis to develop joint business opportunities. After market opportunities and bottlenecks are identified, the stakeholders, organized in thematic groups, will analyze them and develop work plans. For example:

- One thematic group could develop work plans (business plans in this case) for commercial innovations, such as a new brand of certified seed produced and distributed by a farmers' cooperative or a local company.
- Another thematic group could develop work plans to develop technical innovations, such as improved techniques to produce in vitro plants, minitubers or cuttings, or a new diagnostic technique to detect seed-borne diseases, or a seed replacement practice to combine the use of high-quality seed and on-farm practices.
- Another group could develop work plans to develop institutional innovations, such as creating a new regulation for exporting seed to neighboring countries, or for producing quality declared seed, or promoting the creation of a new network of seed multipliers.

The workplans will be presented at a second public event. The intended output is a work plan for each thematic group. This activity will also take approximately three months.

Activity 3.3. Implementation of joint innovations. The final activity is to start the implementation of the work plan by each of the thematic groups. They can request support to conduct specific studies or tasks, for example:

- A business plan for the production and distributions of certified seed by a farmers' cooperative or a local company.
- Experiments to adapt technologies to improve the production of in vitro plants, minitubers or cuttings.
- Experiments to adapt a diagnostic technique to local conditions.
- Experiments to test seed replacement strategies.
- A regulation for exporting seed.

- A regulation for producing quality declared seed.

Consulting firms and experts could be hired to conduct these studies and tasks. The resulting innovations (e.g. a new brand of certified seed, a new seed regulation, a new technique for producing in vitro plants) will be presented at a final public event. This activity will take approximately six months. Follow-up of these activities could take two to five years, depending on the innovation.

References

- Access Agriculture n.d. Cassava videos. Web site available at: <https://www.accessagriculture.org/category/59/cassava> [Accessed September 7, 2018]
- Bernet, T., Thiele, G. and Zschocke, T. 2006. Participatory Market Chain Approach (PMCA) – User Guide. International Potato Center (CIP) – Papa Andina, Lima, Peru. Available at: <http://cipotato.org/wp-content/uploads/2014/09/003296.pdf> [Accessed September 7, 2018]
- Bryan, J. 1983. On-farm seed improvement by the potato seed plot technique. International Potato Center. Lima, Peru. 13 p. Available at: <http://cipotato.org/wp-content/uploads/2014/09/TIBen18058.pdf> [Accessed September 7, 2018]
- Cáceres, P.A., Pumisacho, M., Forbes, G.A. and Andrade-Piedra, J.L. 2008. Learning to Control Potato Late Blight – A Facilitator’s Guide. International Potato Center (CIP), Instituto Nacional Autónomo de Investigaciones Agropecuarias del Ecuador (INIAP), Secretaría Nacional de Ciencia y Tecnología del Ecuador (SENACYT). Quito, Ecuador. 142 p. Available at: <http://cipotato.org/wp-content/uploads/2014/08/004358.pdf> [Accessed September 7, 2018]
- Carli, C., Gulbani, A., Khidesheli, Z. and Kobalatse K. 2010. Strengthening the potato seed system in Georgia: preliminary results. Lima: International Potato Center (CIP): pp. 22. <http://hdl.handle.net/10568/67216>
- De Haan, S., Salas, E., Fonseca, C., Gastelo, M., Amaya, N., Bastos, C., Hualla, V. and Bonierbale, M. 2017. Selección participativa de variedades de papa (SPV) usando el diseño mamá y bebé: una guía para capacitadores con perspectiva de género. Lima (Peru). International Potato Center (CIP). ISBN 978-92-9060-475-4. 82 p. <https://hdl.handle.net/10568/81221>
- Forbes, G.A., Andersen, K., Andrade-Piedra, J.L. Buddenhagen, C., Fulton, J., Gatto, M., Mdivani, R. and Garrett, K. 2018. The potato seed system in the Republic of Georgia: A baseline analysis for a national integrated seed health strategy. Agricultural systems.
- Farm Inputs Promotions Africa (FIPS) n.d. FIPS Promotion of Improved Technology in Kenya through Small Seed- and Fertilizer-Packs. Available at: https://www.future-agricultures.org/wp-content/uploads/pdf-archive/FIPS_SmallPacks_Demos.pdf [Accessed September 7, 2018]

Frost, K.E., Groves, R.L. and Charkowski, A.O. 2013. Integrated control of potato pathogens through seed potato certification and provision of clean seed potatoes. *Plant Dis.* 97:1268–1280.

<https://apsjournals.apsnet.org/doi/pdf/10.1094/PDIS-05-13-0477-FE>

Gatto, M., Hareau, G., Pradel, W., Suarez, V. and Qin, J. 2017. Release and adoption of modern potato varieties in Southeast, East, and South Asia. 9th ASAE International Conference: Transformation in Agricultural and Food Economy in Asia. Bangkok (Thailand). 11-13 Jan 2017. In: 9th ASAE International Conference 2017 Conference Proceedings. Asian Society of Agricultural Economists (ASAE). pp. 1415-1444.

<https://hdl.handle.net/10568/92972>

Geostat. 2019. Food balance sheets. Balance sheet for potato. National Statistical Office for Georgia, T'bilisi, Georgia. (Available from http://www.geostat.ge/index.php?action=page&p_id=752&lang=eng) (Accessed March 11, 2019)

Gildemacher. P., Demo, P., Kinyae, P., Wakahiu, M., Nyongesa, M., Zschocke, T. 2007a. Select the best: positive selection to improve farm saved seed potatoes: Picture book. International Potato Center. Lima, Peru

<https://hdl.handle.net/10568/76796>

Gildemacher. P. Demo, P., Kinyae, P., Wakahiu, M., Nyongesa, M., Zschocke, T. 2007b. Select the best: positive selection to improve farm saved seed potatoes: Farmer field aid. International Potato Center. Lima, Peru.

<https://hdl.handle.net/10568/76795>.

Gildemacher. P. Demo, P., Kinyae, P., Wakahiu, M., Nyongesa, M., Zschocke, T. 2007c. Select the best: positive selection to improve farm saved seed potatoes: Trainers manual. International Potato Center. Lima, Peru.

<https://hdl.handle.net/10568/76797>.


Holzworth, D.P., Huth, N.I., de Voil, P.G., Zurcher, E.J., Herrmann, N.I., McLean, G., Chenu, K. et al. 2014. APSIM - Evolution towards a New Generation of Agricultural Systems Simulation. *Environmental Modelling & Software* 62:327–350. <https://doi.org/10.1016/j.envsoft.2014.07.009>.

International Potato Center (CIP). n.d. Catalogue of CIP advanced clones. Web site available at: <https://research.cip.cgiar.org/redlatinpapa/pages/home.php> [Accessed September 7, 2018]

International Potato Center (CIP). 2006. Procedures for standard evaluation trials of advanced potato clones. An International Cooperators' Guide. Available at:

- <https://research.cip.cgiar.org/confluence/download/attachments/14942262/ICG.pdf?version=1> [Accessed September 7, 2018]
- International Potato Center (CIP). 2016. Research on local seed potato value chain gaps and a bottlenecks in Tajikistan. Potato production support and research to improve food security in Khatlon, Tajikistan. International Potato Center: pp. 19. <https://hdl.handle.net/10568/77152>.
- International Potato Center (CIP). 2018. Enhancing Rural Livelihoods in Georgia: Introducing Integrated Seed Health Approaches to Local Potato Seed Systems. Web site available at: <https://ishpotato.cipotato.org/> [Accessed September 7, 2018]
- Kinyua, Z., et al. 2015. Adaptation and Improvement of the Seed-plot Technique in Smallholder Potato Production. In: J. Low et al. (eds). Potato and Sweetpotato in Africa: Transforming 218 the Value Chains for Food and Nutrition Security. Ninth Triennial African Potato Association Conference, Naivasha, Kenya, 30 June-4 July 2013. Available at: <https://www.cabi.org/cabebooks/ebook/20153367576> [Accessed September 7, 2018]
- Mdivani, R. 2011. [The new potato in high-mountain conditions]. Agrarian Science 11. p. 17. Available at: <http://vetpress.ru/upload/iblock/907/90747d6853eca558407cc7a18ffd0547.pdf> [Accessed September 14, 2018]
- McGuire, S. and Sperling, L. 2016. Seed systems smallholder farmers use. Food Security 8:179–195. doi: 10.1007/s12571-015-0528-8
- Ministry of Agriculture of Georgia (MoA). 2015. Strategy for Agricultural Development in Georgia 2015-2020. Ministry of Agriculture of Georgia: pp. 38. Available at: <http://www.moa.gov.ge/Download/Files/92> [Accessed September 7, 2018]
- Mudege, N.N., Salas, E., De Haan, S., Bonierbale, M., Mendes, T. and Amele, A. 2017. Experience and guidelines for PVS: Integrating gender and potato breeding. Gender and Breeding Meeting. Nairobi, Kenya. 5-6 October 2017. <https://hdl.handle.net/10568/89293>.
- National Statistics Office of Georgia (GeoStat.ge). Web site available at: <http://www.geostat.ge/>
- Obura, B.O., Parker, M.L., Bruns, C., Finckh, M.R. and Schulte-Geldermann, E. 2016. Cost Benefit Analysis of Seed Potato Replacement Strategies among Smallholder Farmers in Kenya. Tropentag, September 18-21, 2016, Vienna, Austria. Available at: http://www.tropentag.de/2016/abstracts/links/Obura_eAhVYqsS.pdf. [Accessed September 7, 2018]

- Ogero, K., McEwan, M. and Pamba, N. Clean vines for smallholder farmers in Tanzania. In: Andrade-Piedra J., Bentley, J., Almekinders, C., Jacobsen, K., Walsh, S., and Thiele, G. (eds.) 2016. Case Studies of Roots, Tubers and Bananas Seed Systems. CGIAR Research Program on Roots, Tubers and Bananas (RTB), Lima: RTB Working Paper No. 2016-3. ISSN 2309-6586. 244 p. <https://hdl.handle.net/10568/81052>.
- Ondity, J., Njoroge, K. and Shibairo, S. 2013. Improving potato tuber yields using genotypes with multiple virus resistance in Kenya. *Agric. Biol. J. N. Am.* 4:406–412. Available at: <https://www.scihub.org/ABJNA/PDF/2013/4/ABJNA-4-4-406-412.pdf> [Accessed September 13, 2018]
- Schulte-Geldermann, E., Gildemacher, P.R. and Struik, P.C. 2012. Improving seed health and seed performance by positive selection in three Kenyan potato varieties. *Am. J. Potato Res.* 89:429–437. doi: 10.1007/s12230-012-9264-1
- Thiele, G., Devaux, A., Reinoso, I., Pico, H., Montesdeoca, F., Pumisacho, M., Andrade-Piedra, J., Velasco, C., Flores, P., Esprella, R., Thomann, A., Manrique, K., and Horton, D. 2011. Multi-stakeholder platforms for linking small farmers to value chains: evidence from the Andes. *Int. J. Agric. Sustain.* 9:423–433. <https://doi.org/10.1080/14735903.2011.589206>
- Thomas-Sharma, S., Abdurahman, A., Ali, S., Andrade-Piedra, J. L., Bao, S., Charkowski, A. O., et al. 2015. Seed degeneration in potato: The need for an integrated seed health strategy to mitigate the problem in developing countries. *Plant Pathol.* 65:3–16. doi: 10.1111/ppa.12439
- Wahyuning, S. et al. 2006. Farmer Field School for Potato Integrated Pest Management A Facilitator’s Field Guide. International Potato Center and FAO Regional Vegetable IPM Program in South and Southeast Asia. Available at: <http://www.vegetableipmasia.org/uploads/files/document/TrainingMaterials/Potato-IPM-FFS-exercise-manual.pdf> [Accessed September 7, 2018]



CIP is a research-for-development organization with a focus on potato, sweetpotato and Andean roots and tubers. It delivers innovative science-based solutions to enhance access to affordable nutritious food, foster inclusive sustainable business and employment growth, and drive the climate resilience of root and tuber agri-food systems. Headquartered in Lima, Peru, CIP has a research presence in more than 20 countries in Africa, Asia and Latin America.

www.cipotato.org

CIP is a CGIAR research center

CGIAR is a global research partnership for a food-secure future. Its science is carried out by 15 research centers in close collaboration with hundreds of partners across the globe.

www.cgiar.org

For more information, please contact CIP Headquarter. Av. La Molina 1895, La Molina. Apartado 1558, Lima 12, Peru.

 5-11-3496017  cip-cpad@cgiar.org  www.cipotato.org |  @cipotato  @Cipotato  @cip_cipotato

CIP thanks all donors and organizations which globally support its work through their contributions to the CGIAR Trust Fund.

<https://www.cgiar.org/funders/>



© 2019. International Potato Center. All rights reserved.

This work by the International Potato Center is licensed under a Creative Commons Attribution 4.0 International (CC BY 4.0). To view a copy of this license, visit <https://creativecommons.org/licenses/by/4.0/>. Permissions beyond the scope of this license may be available at: <http://www.cipotato.org/contact/>